KIAS-P12065

Two-loop effective potential, thermal resummation and first-order phase transitions: Beyond high-temperature expansion

Koichi Funakubo[1]* and Eibun Senaha[2]†
[1] *Department of Physics, Saga University, Saga 840-8502 Japan and*
[2] *School of Physics, KIAS, Seoul 130-722, Korea*

(Dated: October 8, 2012)

We formulate a tractable calculation scheme for a finite temperature two-loop resummed effective potential in gauge theories without using a high-temperature expansion. We apply it to the Abelian-Higgs model and its extension to a minimal supersymmetric standard model-like model and study the thermal phase transition. It is shown that our scheme improves the previous results about 10% in the quantities relevant to the phase transition, and its impacts on bubble dynamics could be even more sizable. It still holds that the stop-stop-gluon sunset diagram enhances the strength of the first-order phase transition even without the high-temperature expansion.

PACS numbers: 05.70.Fh, 11.10.Wx, 12.38.Cy, 98.80.Cq

arXiv:1210.1737v1 [hep-ph] 5 Oct 2012

Introduction. The finite-temperature (T) effective potential is a powerful tool for studying various phenomena in hot early Universe, for example, thermal phase transitions at quantum chromodynamics (QCD) and electroweak scales. It is known that a first-order electroweak phase transition (EWPT) is one of the necessary conditions for successful electroweak baryogenesis (EWBG) [1]. A naive perturbation method at high T is known as unreliable. Generally, the perturbative expansion in a coupling constant is invalidated at high T due to the sizable T-dependent loop corrections. Therefore, such higher-order corrections should be resummed to obtain a sensible result. In the EWBG context, the EWPT up to two-loop order have been studied so far. However, since the calculation of the resummed two-loop effective potential is notoriously a formidable task, a high-T expansion (HTE) in which a particle mass is less than temperature was exclusively used. Since the relevant parameter region for the successful EWBG is $v_C/T_C \gtrsim 1$, where T_C denotes a critical temperature and v_C is a Higgs vacuum expectation value (VEV) at T_C, the validity of the HTE is far obvious. Indeed, it turns out that the HTE of the two-loop sunset diagrams is valid only for a small mass ratio to temperature, < 0.01 [2], rendering all existing two-loop order results of the EWPT based on the HTE in literature less precise.

In the minimal supersymmetric standard model (MSSM), it is known that v_C/T_C at the one-loop order is not large enough to satisfy the sphaleron decoupling condition even at a bubble nucleation temperature [3]. However, the deficit in v_C/T_C can be compensated by a potentially sizable two-loop sunset diagram consisting of scalar top (stop) and gluon [4, 5]. Therefore, it is of great importance that such a correction is quantified on more stuff basis than the HTE. The viability of the MSSM baryogenesis depends not only on the current Large Hadron Collider data but potentially on the two-loop calculation of the EWPT beyond the HTE. For the former, refer to the recent papers [6].

In this Letter, we formulate a tractable calculation scheme for the two-loop effective potential and thermal resummation without relying on the HTE. In order to go beyond the HTE, in which T-dependent divergent terms are omitted, we must execute the renormalization with T-dependent counter terms arising from the resummation. As an example, we demonstrate our calculation scheme in the Abelian-Higgs (AH) model and apply it to a MSSM-like model for studying the thermal PT.

Model. Let us begin by introducing the AH model and define the various notations used here. The Lagrangian of the AH is

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + |D_\mu\Phi|^2 - V_0(|\Phi|^2), \tag{1}$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, $D_\mu\Phi = (\partial_\mu - ieA_\mu)\Phi$ and The scalar potential is given by

$$V_0(|\Phi|^2) = -\nu^2|\Phi|^2 + \frac{\lambda}{4}|\Phi|^4. \tag{2}$$

We parametrize the scalar field in terms of the VEV (v) and fluctuation fields

$$\Phi = \frac{1}{\sqrt{2}}(v + h + ia), \tag{3}$$

where h is a physical state and a is a Nambu-Goldstone (NG) boson which is eaten by the gauge boson. The field-dependent scalar and gauge boson masses are

$$m_h^2 = -\nu^2 + \frac{3\lambda}{4}v^2, \quad m_a^2 = -\nu^2 + \frac{\lambda}{4}v^2, \quad m_A^2 = e^2v^2, \tag{4}$$

where we work in the Landau gauge. Throughout the paper, we will adopt $\overline{\text{MS}}$ scheme for renormalization.

Resummation method. We closely follow the resummation method presented in Ref. [7]. The basic procedure of this resummation is simply to add and subtract the T-dependent mass terms $\Delta_T m^2$ in the original Lagrangian.

First, we focus on the mass term of h. The resummed Lagrangian and new counter terms in the d dimension are respectively given by

$$V_0 = \frac{1}{2}\left(-\nu^2 + \Delta_T m_h^2 + \frac{3\lambda\mu^\epsilon}{4}v^2\right)h^2 + \dots, \tag{5}$$

$$\delta_T V = \frac{1}{2}\left(-\delta\nu^2 - \Delta_T m_h^2 + \frac{3\delta\lambda\mu^\epsilon}{4}v^2\right)h^2 + \dots, \tag{6}$$

where $\epsilon = 4-d$, $\delta\nu^2$ and $\delta\lambda$ are the usual $\overline{\rm MS}$ counter terms that are determined by the $1/\epsilon$ terms, and μ is a renormalization scale. The same procedure holds for the mass of a. With the resummed Lagrangian, the scalar propagators are

$$\Delta_h(p) = \frac{1}{p^2 - m_h^2(T)}, \quad \Delta_a(p) = \frac{1}{p^2 - m_a^2(T)}, \tag{7}$$

where $m_{h,a}^2(T) = m_{h,a}^2 + \Delta_T m_{h,a}^2$.

The above resummation scheme has been applied to various models with and without spontaneous symmetry breaking. The proofs of renormalizability and NG theorem in $O(N)$ ϕ^4 theory are given in Ref. [8].

For the gauge sector, similarly, we add and subtract a thermal correction in the original Lagrangian. In this case, however, we should treat longitudinal and transverse parts of the gauge boson propagator separately since thermal corrections to them are different from each other. The resummed bare Lagrangian is given by

$$\begin{aligned}\mathcal{L}_B = \mathcal{L}_B^{\rm unresum} &+ \frac{1}{2}A^\mu\Big[\Delta_T m_L^2 L_{\mu\nu}(i\partial) + \Delta_T m_T^2 T_{\mu\nu}(i\partial)\Big]A^\nu \\ &- \frac{1}{2}A^\mu\Big[\Delta_T m_L^2 L_{\mu\nu}(i\partial) + \Delta_T m_T^2 T_{\mu\nu}(i\partial)\Big]A^\nu,\end{aligned} \tag{8}$$

where $\mathcal{L}_B^{\rm unresum}$ is the unresummed bare Lagrangian and $L_{\mu\nu}(p)$ and $T_{\mu\nu}(p)$ are the projection tensors which take the form of

$$T_{00} = T_{0i} = T_{i0} = 0, \quad T_{ij} = g_{ij} - \frac{p_i p_j}{-\boldsymbol{p}^2}, \tag{9}$$

$$L_{\mu\nu} = P_{\mu\nu} - T_{\mu\nu}, \quad P_{\mu\nu} = g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2}, \tag{10}$$

in the rest frame of the thermal bath. Note that additional T-dependent terms are nonlocal and noncovariant, but gauge invariance is still maintained. With Eq. (8), the resummed gauge boson propagator is cast into the form

$$\begin{aligned}\boldsymbol{D}_{\mu\nu}(p) &= \frac{-1}{p^2 - m_L^2}L_{\mu\nu}(p) + \frac{-1}{p^2 - m_T^2}T_{\mu\nu}(p) \\ &= \left[\frac{-(1-r)}{p^2 - m_L^2} + \frac{-r}{p^2 - m_T^2}\right]P_{\mu\nu}(p) \\ &\quad + \left[\frac{-1}{p^2 - m_T^2} - \frac{-1}{p^2 - m_L^2}\right]\big(T_{\mu\nu}(p) - rP_{\mu\nu}(p)\big),\end{aligned} \tag{11}$$



(a) (b) (c) (d) (CT)

FIG. 1. The diagrams in which the gauge boson is involved in the AH model.



(a) (b) (c) (d)

FIG. 2. The IPI two-point functions of A_μ.

where $m_{L,T}^2 = m_A^2 + \Delta_T m_{L,T}^2$ and r is an arbitrary real parameter. Note that the noncovariant part in the last line of Eq. (11) yields less ultra-violet divergent loop integrals. Requiring $g^{\mu\nu}\big(T_{\mu\nu}(p) - rP_{\mu\nu}(p)\big) = 0$, one gets $r = (d-2)/(d-1)$. We will see that this specific choice of r makes it easy to perform the two-loop calculations significantly. Let us denote the second and third lines in Eq. (11) by $\boldsymbol{D}_{\mu\nu}(p) = D_{\mu\nu}^{\rm cov}(p) + \delta D_{\mu\nu}(p)$. With this gauge boson propagator, together with Eq. (7), we will compute the two-loop diagrams. Since there is no difficulty to obtain the resummed diagrams for purely scalar sector, we here concentrate on the diagrams involving the gauge boson as depicted in FIG. 1. Each diagram is expressed as

$$\begin{aligned}\mu^\epsilon V_R^{\rm (a)}(v;T) = \frac{e^2}{2}\sum\!\!\!\!\!\!\!\int_k \sum\!\!\!\!\!\!\!\int_q (2k+q)^\mu(2k+q)^\nu \\ \times \Delta_h(k)\Delta_a(k+q)\boldsymbol{D}_{\mu\nu}(q),\end{aligned} \tag{12}$$

$$\mu^\epsilon V_R^{\rm (b)}(v;T) = e^2\mu^\epsilon v^2 \sum\!\!\!\!\!\!\!\int_k \sum\!\!\!\!\!\!\!\int_q \Delta_h(k+q)\boldsymbol{D}^{\mu\nu}(k)\boldsymbol{D}_{\mu\nu}(q), \tag{13}$$

$$\mu^\epsilon V_R^{\rm (c)}(v;T) = -\frac{1}{2}e^2 g^{\mu\nu}\sum\!\!\!\!\!\!\!\int_k \sum\!\!\!\!\!\!\!\int_q \Delta_h(k)\boldsymbol{D}_{\mu\nu}(q), \tag{14}$$

$$\mu^\epsilon V_R^{\rm (d)}(v;T) = -\frac{1}{2}e^2 g^{\mu\nu}\sum\!\!\!\!\!\!\!\int_k \sum\!\!\!\!\!\!\!\int_q \Delta_a(k)\boldsymbol{D}_{\mu\nu}(q), \tag{15}$$

$$\begin{aligned}\mu^\epsilon V_R^{\rm (CT)}(v;T) = \frac{1}{2}\sum\!\!\!\!\!\!\!\int_q \Big(-\hat{\Gamma}_A(q)\Big|_{\rm div} - \Delta_T m_L^2 L(q) \\ - \Delta_T m_T^2 T(q)\Big)^{\mu\nu}\boldsymbol{D}_{\mu\nu}(q),\end{aligned} \tag{16}$$

where the sum-integral symbol is defined by

$$\sum\!\!\!\!\!\!\!\int_k = \mu^\epsilon T\sum_{n=-\infty}^{\infty}\int \frac{d^{d-1}\boldsymbol{k}}{(2\pi)^{d-1}}, \tag{17}$$

with $k^0 = k_0 = i\omega_n = 2n\pi iT$ and $\hat{\Gamma}_A^{\mu\nu}(q)$ is the one-particle-irreducible (1PI) two-point function of A_μ evaluated with the propagator $D_{\mu\nu}^{\rm cov}$. Now we expand above $\mu^\epsilon V_R(v;T)$'s in powers of $\delta D_{\mu\nu}$ as

$$\begin{aligned}
&\mu^\epsilon\Big(V_R^{\rm (a)}+V_R^{\rm (b)}+V_R^{\rm (c)}+V_R^{\rm (d)}+V_R^{\rm (CT)}\Big)[\boldsymbol{D}]\\
&\quad=\mu^\epsilon\Big(V_R^{\rm (a)}+V_R^{\rm (b)}+V_R^{\rm (c)}+V_R^{\rm (d)}+V_R^{\rm (CT)}\Big)[D^{\rm cov}]\\
&\qquad+\frac{1}{2}\sum\!\!\!\!\!\!\!\int_q\Big[\hat{\Gamma}_A^{\rm (a)}(q)+\hat{\Gamma}_A^{\rm (b)}(q)+\hat{\Gamma}_A^{\rm (c)}(q)+\hat{\Gamma}_A^{\rm (d)}(q)-\hat{\Gamma}_A(q)\Big|_{\rm div}\\
&\qquad\qquad-\Delta_T m_L^2 L(q)-\Delta_T m_T^2 T(q)\Big]^{\mu\nu}\delta D_{\mu\nu}(q)\\
&\qquad+e^2\mu^\epsilon v^2\sum\!\!\!\!\!\!\!\int_k\sum\!\!\!\!\!\!\!\int_q\Delta_h(k+q)\delta D^{\mu\nu}(k)\delta D_{\mu\nu}(q). \qquad (18)
\end{aligned}$$

Note that the expanded terms to first order in derivative can be written in terms of $\hat{\Gamma}_A^i(q)$ with $i=(a)-(d)$ which are represented in FIG. 2. For the covariant sector in the first line, one can show that all T-dependent divergences are cancelled out and renormalization is successfully carried out in the ordinary manner [9]. Hence, we here prove finiteness of the noncovariant sector.

In general, because of the gauge invariance the polarization tensor is written as

$$\begin{aligned}
\Pi_{\mu\nu}(q)&=\Pi_0(q^2)P_{\mu\nu}(q)\\
&\quad+\Delta\Pi_L(q^0,\boldsymbol{q})L_{\mu\nu}(q)+\Delta\Pi_T(q^0,\boldsymbol{q})T_{\mu\nu}(q). \qquad (19)
\end{aligned}$$

$\Delta\Pi_{L,T}(q^0,\boldsymbol{q})$ in the infrared (IR) limit ($q^0=0$ and $\boldsymbol{q}\to 0$) may take the form

$$\Delta\Pi_{L,T}(q^0,\boldsymbol{q})\to\Delta_T m_{L,T}^2+\frac{\epsilon}{2}\pi_{L,T}^{(\epsilon)}(T), \qquad (20)$$

Thus, in the IR limit the second and third lines of Eq. (18) are reduced to

$$\begin{aligned}
&\frac{1}{2}\sum\!\!\!\!\!\!\!\int_q\Big[\hat{\Pi}(q)-\Delta_T m_L^2 L(q)-\Delta_T m_T^2 T(q)\Big]^{\mu\nu}\delta D_{\mu\nu}(q)\\
&\quad=\frac{m_L^2-m_T^2}{48\pi^2}\Big(\pi_T^{(\epsilon)}(T)-\pi_L^{(\epsilon)}(T)\Big)+\mathcal{O}(\epsilon). \qquad (21)
\end{aligned}$$

Therefore, we obtain the finite result. Note that the subtraction terms $\Delta_T m_{L,T}^2$ in the counter term play an important role in this proof. It is easy to show the finiteness of this noncovariant part without the IR limit as well [9]. Likewise, the last line of Eq. (18) is also finite [9]. It should be stressed that so far the HTE is not used.

Before studying the PT, we look into some approximations that are commonly used in the finite-T two-loop calculations. Let us define

$$\begin{aligned}
&H(m_1,m_2,m_3)\\
&\quad=\sum\!\!\!\!\!\!\!\int_k\sum\!\!\!\!\!\!\!\int_q\frac{1}{(k^2+m_1^2)(q^2+m_2^2)\big[(k+q)^2+m_3^2\big]},\\
&\qquad+\frac{T^2}{4(2\pi)^4}\sum_{\{i,j,k\}=\text{cyclic}}^{1,2,3}K_{--}(a_i,a_j,a_k)+\ldots, \qquad (22)
\end{aligned}$$



FIG. 3. $K_{--}(a,a,0)$, $3K(2a/3)$ and $3K^{\rm HTE}(2a/3)$ are plotted as a function of a.

where $a_i = m_i/T$ and the momentum is defined in the Euclidean space, and so $k^2=\omega_m^2+\boldsymbol{k}$, $q^2=\omega_n^2+\boldsymbol{q}^2$, *etc.* $K_{--}(a_i,a_j,a_k)$ is defined by the function that contains two $n_B(E)$, where, for instance, $n_B(E_k)=1/(e^{E_k/T}-1)$ with $E_k=\sqrt{\boldsymbol{k}^2+m_1^2}$. We define $K_{--}(a,a,a)=K(a)$ [7]. The ellipses stand for the zero-T contributions and the terms including one $n_B(E)$, which can be expressed in terms of the one-loop finite-T integrals.

In literature, in addition to the HTE, the following approximation is frequently used [10]

$$H(m_1,m_2,m_3)\to H\left(\frac{m_1+m_2+m_3}{3}\right). \qquad (23)$$

We investigate the validity of this mass-averaging approximation without the HTE. Here, we take $a_1=a_2\equiv a$ and $a_3=0$. This particular choice corresponds to, for example, the stop-stop-gluon sunset diagram in the MSSM. The numerical comparison of $K_{--}(a,a,0)$, $3K(2a/3)$ and $3K^{\rm HTE}(2a/3)$ [7] are shown in FIG. 3. We can see that all of them would agree in the $a\to 0$ limit. It is found that $K_{--}(a,a,0)/(3K(2a/3))\lesssim 0.6$ for $a\gtrsim 1$. However, the corresponding value of each function is less than one. So the error of the mass-averaging approximation can be small [9]. On the contrary, the use of the HTE gives a large deviation and yet negative value at $a=\mathcal{O}(1)$ even though $K(a)$ is positive-definite by definition. It is concluded that the HTE of $K(a)$ is valid only for $a\lesssim 0.01$, which is consistent with the results in Ref. [2].

Application to a MSSM-like model. Now we move on to discuss the thermal PT using the scheme formulated in this Letter. As a first step toward the complete analysis of the two-loop driven first-order EWPT scenario such as the MSSM, we consider an extended AH model in which additional $U(1)$ gauge boson and complex scalar



FIG. 4. v_C/T_C in the three cases are shown as a function of g_3. The input parameters are given in the text.



FIG. 5. The shown is the effective potential at T_C. We take $g_3 = 1.2$. The remaining parameters are the same as in FIG. 4.

are introduced. The added Lagrangian is

$$\Delta\mathcal{L} = -\frac{1}{4}G_{\mu\nu}G^{\mu\nu} + |D_\mu\tilde{t}|^2 - (m_0^2 + y^2|\Phi|^2)|\tilde{t}|^2 + \frac{\tilde{\lambda}}{4}|\tilde{t}|^4, \tag{24}$$

where $G_{\mu\nu} = \partial_\mu G_\nu - \partial_\nu G_\mu$ and $D_\mu\tilde{t} = (\partial_\mu - ig_3G_\mu)\tilde{t}$. As is done in the AH model, we carry out the thermal resummation in our scheme. To be specific, we keep only $\mathcal{O}(T^2)$ corrections in $\Delta_T m^2$ and $\pi^{(\epsilon)}(T)$.

As mentioned above, in the MSSM the stop-stop-gluon sunset diagram enhances v_C/T_C. We scrutinize this effect with and without the HTE. In what follows, by the HTE we mean that only K_{--} is evaluated using the HTE.

In FIG. 4, v_C/T_C is shown as a function of g_3. We set $v = 246$, $m_h = 35$, $m_0^2 = 0$, $y = 1.0$, $e = 0.5$, $\tilde{\lambda} = 0.3$ and $\bar{\mu} = 2\sqrt{\pi}e^{-\gamma_E/2}\mu = 150$, where $\gamma_E \simeq 0.577$. The red curve represents the two-loop calculation, and the blue dashed curve denotes the two-loop calculation with the HTE. The one-loop calculation of v_C/T_C is also shown by the dotted black curve. This figure shows that enhancement of v_C/T_C due to the $\tilde{t}$-$\tilde{t}$-G_μ diagram can still persist beyond the HTE. In this specific example, the use of the HTE leads to the underestimated v_C/T_C.

The height of the barrier between the two degenerate vacua in the effective potential (denoted by h_b) is also relevant to dynamics of the first-order PT. The effective potentials at T_C and $g_3 = 1.2$ in the three cases are plotted in FIG. 5. The color and line coordinates are the same as in FIG. 4. We find that $v_C/T_C|_{\text{1-loop}} = 186.55/76.75 = 2.43$, $v_C/T_C|_{\text{2-loop}} = 191.80/74.81 = 2.56$ and $v_C/T_C|_{\text{2-loop HTE}} = 204.94/81.32 = 2.52$. The significant increase of T_C in the HTE case may be the consequence of the artificial negative contributions to the quadratic term in the scalar potential [9]. It is also found that h_b at the two-loop level is somewhat higher than that of the one-loop case, delaying the onset of the PT. However, we may get the overestimated result once the HTE is used. We observe that generally the larger g_3 can bring the larger errors in v_C, T_C and h_b.

Conclusion. We have formulated the tractable calculation scheme for the finite-T two-loop resummed effective potential in gauge theories. Our analysis on the PT indicates that for the typical parameter set the use of the HTE in the sunset diagrams can lead to the overestimated results by about 10% in v_C and T_C and by around 50% in h_b. Since our calculation scheme is model-independent, it is applicable for any $U(1)$ models. Even in the non-Abelian case, we expect that the dominant effects relevant to the PT would be modeled by the sunset diagrams in the AH or its extended models since a gluon-gluon-gluon type sunset diagram has little effect.

---

* funakubo@cc.saga-u.ac.jp
† senaha@kias.re.kr

[1] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **155**, 36 (1985).
[2] M. Laine and M. Losada, Nucl. Phys. B **582**, 277 (2000).
[3] K. Funakubo and E. Senaha, Phys. Rev. D **79**, 115024 (2009).
[4] J. R. Espinosa, Nucl. Phys. B **475**, 273 (1996).
[5] M. Carena, G. Nardini, M. Quiros and C. E. M. Wagner, Nucl. Phys. B **812**, 243 (2009).
[6] T. Cohen, D. E. Morrissey and A. Pierce, Phys. Rev. D **86**, 013009 (2012); D. Curtin, P. Jaiswal and P. Meade, JHEP **1208**, 005 (2012); M. Carena, G. Nardini, M. Quiros and C. E. M. Wagner, arXiv:1207.6330 [hep-ph].
[7] R. R. Parwani, Phys. Rev. D **45**, 4695 (1992) [Erratum-ibid. D **48**, 5965 (1993)].
[8] S. Chiku and T. Hatsuda, Phys. Rev. D **58**, 076001 (1998).
[9] K. Funakubo and E. Senaha, in preparation.
[10] P. B. Arnold and O. Espinosa, Phys. Rev. D **47**, 3546 (1993) [Erratum-ibid. D **50**, 6662 (1994)].